



## Human Epidermal Melanocytes-dark (HEM-d)

Catalog #2220 (old Cat. #2200)

### Cell Specification

Melanocytes are neural crest-derived cells that produce melanin via melanogenesis. Melanocytes localize to several tissues including the epidermis, eye, inner ear and leptomeninges. Dysregulation of melanocyte migration, proliferation, or survival during embryonic development thus causes congenital disorders in those tissues as seen in Tietz syndrome, Waardenburg syndrome, and piebaldism [1]. In the bottom layer of skin epidermis, melanocytes synthesize and transfer dark-colored melanin to surrounding keratinocytes to give skin pigmentation. Melanin also blocks UV-B light to protect the hypodermis from solar exposure-induced photodamage. Progress in culture techniques, along with an improved understanding of melanocyte biology, has led to a successful culture system to model melanomas, inner ear homeostasis, vitiligo, and mitochondrial dysfunction in Duchenne Muscular Dystrophy [2-4].

HEM-d from ScienCell Research Laboratories is isolated from neonatal human skin. HEM-d are cryopreserved at passage one and delivered frozen. Each vial contains  $>5 \times 10^5$  cells in 1 ml volume. HEM-d are characterized by immunofluorescence with antibodies specific to fibronectin and NGF-receptor (p75). HEM-d are negative for HIV-1, HBV, HCV, mycoplasma, bacteria, yeast, and fungi. HEM-d are guaranteed to further expand for 10 population doublings under the conditions provided by ScienCell Research Laboratories.

### Recommended Medium

It is recommended to use Melanocyte Medium (MeLM, Cat. #2201) for culturing HEM-d *in vitro*.

### Product Use

HEM-d are for research use only. They are not approved for human or animal use, or for application in *in vitro* diagnostic procedures.

### Storage

Upon receiving, directly and immediately transfer the cells from dry ice to liquid nitrogen and keep the cells in liquid nitrogen until they are needed for experiments.

### Shipping

Dry ice.

### References

- [1] Li J, Song JS, Bell RJ, Tran TN, Haq R, Liu H, Love KT, Langer R, Anderson DG, Larue L, Fisher DE. (2012) "YY1 regulates melanocyte development and function by cooperating with MITF." *PLoS Genet.* 8: e1002688.
- [2] Neng L, Zhang F, Kachelmeier A, Shi X. (2013) "Endothelial cell, pericyte, and perivascular resident macrophage-type melanocyte interactions regulate cochlear intrastrial fluid-blood barrier permeability." *J Assoc Res Otolaryngol.* 14: 175-85.
- [3] Ellassiuty YE, Klarquist J, Speiser J, Yousef RM, El Refaee AA, Hunter NS, Shaker OG, Gundeti M, Nieuweboer-Krobotova L, Le Poole IC. (2011) "Heme oxygenase-1 expression protects melanocytes from stress-induced cell death: implications for vitiligo." *Exp Dermatol.* 20: 496-501.
- [4] Pellegrini C, Zulian A, Gualandi F, Manzati E, Merlini L, Michelini ME, Benassi L, Marmioli S, Ferlini A, Sabatelli P, Bernardi P, Maraldi NM. (2013) "Melanocytes--a novel tool to study mitochondrial dysfunction in Duchenne muscular dystrophy." *J Cell Physiol.* 228: 1323-31.

## Instructions for culturing primary cells

---

**Caution:** Cryopreserved primary cells are very delicate. Thaw the vial in a 37°C water bath and return the cells to culture as quickly as possible with minimal handling! Do not centrifuge the cells after thawing as this can damage the cells.

### Initiating the culture:

**Note:** ScienCell primary cells must be cultured in a 37°C, 5% CO<sub>2</sub> incubator. Cells are only warranted if ScienCell media and reagents are used and the recommended protocols are followed.

1. Prepare a poly-L-lysine-coated culture vessel (2 µg/cm<sup>2</sup>, T-75 flask is recommended). To obtain a 2 µg/cm<sup>2</sup> poly-L-lysine-coated culture vessel, add 10 ml of sterile water to a T-75 flask and then add 15 µl of poly-L-lysine stock solution (10 mg/ml, Cat. #0413). Leave the vessel in a 37°C incubator overnight (or for a minimum of one hour).
2. Prepare complete medium. Decontaminate the external surfaces of medium bottle and medium supplement tubes with 70% ethanol and transfer them to a sterile field. Aseptically transfer supplement to the basal medium with a pipette. Rinse the supplement tube with medium to recover the entire volume.
3. Rinse the poly-L-lysine-coated vessel twice with sterile water and then add 15 ml of complete medium. Leave the vessel in the sterile field and proceed to thaw the cryopreserved cells.
4. Place the frozen vial in a 37°C water bath. Hold and rotate the vial gently until the contents completely thaw. Promptly remove the vial from the water bath, wipe it down with 70% ethanol, and transfer it to the sterile field.
5. Carefully remove the cap without touching the interior threads. Gently resuspend and dispense the contents of the vial into the equilibrated, poly-L-lysine-coated culture vessel.

***Note: Dilution and centrifugation of cells after thawing are not recommended since these actions are more harmful to the cells than the effect of residual DMSO in the culture. It is also important that cells are plated in poly-L-lysine-coated culture vessels to promote cell attachment.***

6. Replace the cap or lid of the culture vessel and gently rock the vessel to distribute the cells evenly. Loosen cap, if necessary, to allow gas exchange.
7. Return the culture vessel to the incubator.
8. Do not disturb the culture for at least 16 hours after initiation. Refresh culture medium the next day to remove residual DMSO and unattached cells.

### Maintaining the culture:

1. Refresh supplemented culture medium the next morning after establishing a culture from cryopreserved cells.
2. Change the medium every three days, until the culture is approximately 70% confluent.
3. Once the culture reaches 70% confluency, change medium every other day until the culture is approximately 90% confluent.

## Subculturing:

1. Subculture when the culture reaches 90% confluency.
2. Prepare poly-L-lysine-coated culture vessels ( $2 \mu\text{g}/\text{cm}^2$ ) one day before subculture.
3. Warm complete medium, trypsin/EDTA solution, 0.05% (T/E, Cat. #0183), T/E neutralization solution (TNS, Cat. #0113), and DPBS ( $\text{Ca}^{++}$ - and  $\text{Mg}^{++}$ -free, Cat. #0303) to **room temperature**. We do not recommend warming reagents and medium in a  $37^\circ\text{C}$  water bath prior to use.
4. Rinse the cells with DPBS.
5. Add 5 ml DPBS and 5 ml 0.05% T/E solution (Cat. #0183) into flask (in the case of a T-75 flask). Gently rock the flask to ensure complete coverage of cells by T/E solution. Use a microscope to monitor the change in cell morphology.

**Note:** We recommend using ScienCell 0.05% T/E solution which is optimized to minimize cell damage due to over trypsinization. If 0.25% T/E solution (Cat. #0103) is used, then 9 ml of DPBS and 1 ml of 0.25% T/E solution should be used.

**Caution:** Do NOT use undiluted trypsin when subculturing primary cells.

6. During incubation, prepare a 50 ml conical centrifuge tube with 5 ml of fetal bovine serum (FBS, Cat. #0500).
7. Once cells completely round up, transfer T/E solution from the flask to the 50 ml centrifuge tube (a small percent of cells may detach) and continue to incubate the flask at  $37^\circ\text{C}$  for another minute (no solution in the flask at this time).
8. At the end of incubation, gently tap the side of the flask to dislodge cells from the surface. Check under a microscope to make sure that all cells detach.
9. Add 5 ml of TNS solution to the flask and transfer detached cells to the 50 ml centrifuge tube. Rinse the flask with another 5 ml of TNS to collect the residual cells.
10. Examine the flask under a microscope for a successful cell harvest by looking at the number of cells being left behind; there should be less than 5%.
11. Centrifuge the 50 ml centrifuge tube at 1000 rpm for 5 minutes. Gently resuspend cells in culture medium.
12. Count and plate cells in a new poly-L-lysine-coated culture vessel with the recommended cell density. A seeding density of  $5,000 \text{ cells}/\text{cm}^2$  is recommended.

**Note:** We do not recommend cryopreservation of primary cells by the end user. Refreezing cells may damage them and affect cell performance. ScienCell does not guarantee primary cells cryopreserved by the end user.

*Caution: Handling human derived products is potentially biohazardous. Although each cell strain tests negative for HIV, HBV and HCV DNA, diagnostic tests are not necessarily 100% accurate, therefore, proper precautions must be taken to avoid inadvertent exposure. Always wear gloves and safety glasses when working with these materials. Never mouth pipette. We recommend following the universal procedures for handling products of human origin as the minimum precaution against contamination [1].*

[1] Grizzle WE, Polt S. (1988) "Guidelines to avoid personal contamination by infective agents in research laboratories that use human tissues." *J Tissue Cult Methods*. 11: 191-9.