

# Prostate Epithelial Cell Medium (PEpiCM)

Catalog #4411

### **Product Description**

Prostate Epithelial Cell Medium (PEpiCM), when used with Prostate Epithelial Cell Growth Supplement (PEpiCGS, Cat #4452), is a complete medium designed for optimal growth of normal prostate epithelial cells *in vitro*. It is a sterile, liquid medium which contains essential and non-essential amino acids, vitamins, organic and inorganic compounds, hormones, growth factors and trace minerals. The medium is HEPES and bicarbonate buffered and has a pH of 7.4 when equilibrated in an incubator with an atmosphere of 5% CO<sub>2</sub>/95% air. The medium is formulated (quantitatively and qualitatively) to provide an optimally balanced nutritional environment that selectively supports the growth of normal prostate epithelial cells *in vitro*.

## **Components**

PEpiCM consists of 500 ml of basal medium, 5 ml of Prostate Epithelial Cell Growth Supplement (PEpiCGS, Cat. #4452) and 5 ml of penicillin/streptomycin solution (P/S, Cat. #0503). *Note: PEpiCGS and P/S are not pre-mixed in PEpiCM; they must be added separately to make the complete PEpiCM.* 

<u>Note:</u> Due to bovine pituitary extract (BPE) in the growth supplement, the color may vary with different lots and formation of lipoproteins can result in precipitates. This does not affect the biological activity.

#### **Product Use**

PEpiCM is for research use only. It is not approved for human or animal use, or for application in *in vitro* diagnostic procedures.

#### **Storage**

Store the basal medium at 4°C and the PEpiCGS and P/S solution at -20°C. Protect from light.

#### Shipping

Basal medium: room temperature. Supplements: dry ice.

#### **Instructions for Use**

Thaw PEpiCGS and P/S solution at 37°C. Gently tilt the tubes several times to ensure complete mixing. (**Note**: the presence of BPE in the growth supplement may result in lipoprotein precipitates; this does **not** reduce the efficacy of the supplement). Spray the medium bottle and tubes with 70% ethanol, and wipe to remove excess liquid. In a sterile field, remove the caps without touching the interior threads with fingers. Add PEpiCGS and P/S to the medium and mix well. Since several components are light-labile, the medium should not be exposed to light for extended periods. We do not recommend warming medium in a 37°C water bath prior to use. When stored in the dark at 4°C, the reconstituted medium is stable for one month.

# Rev. 1

Caution: If handled improperly, some components of the medium may present a health hazard. Take appropriate precautions when handling it, including the wearing of protective clothing and eyewear. Dispose of properly.